arXiv:1207.4542v1 [hep-th] 19 Jul 2012

Tricritical gravity waves in the four-dimensional generalized massive gravity

Taeyoon Moon[a*] and Yun Soo Myung[b†],

[a] Center for Quantum Space-time, Sogang University, Seoul, 121-742, Korea
[b] Institute of Basic Sciences and School of Computer Aided Science, Inje University
Gimhae 621-749, Korea

Abstract

We construct a generalized massive gravity by combining quadratic gravity with the Chern-Simons modified term in four dimensions. This is a parity-odd gravity theory. Considering the AdS$_4$ vacuum solution, we derive the linearized Einstein equation, which is not similar to that of the three dimensional generalized massive gravity. For the Kerr-Schild form of a perturbed metric tensor, the linearized equation reduces to a single massive scalar equation. At the tricritical points, we obtain a log-square wave solution. This is compared clearly to the three-dimensional tricritical generalized massive gravity whose dual field theory is a rank-3 logarithmic conformal field theory.

PACS numbers:

Typeset Using LaTeX

---

[*] e-mail address: tymoon@sogang.ac.kr
[†] e-mail address: ysmyung@inje.ac.kr

1 Introduction

The search for a consistent quantum gravity is mainly being suffered from obtaining a renormalizable and unitary quantum field theory. Stelle has first introduced curvature squared terms of $a(R_{\mu\nu}^2 - R^2/3) + bR^2$ in addition to the Einstein-Hilbert term of R [1]. If $ab \neq 0$, the renormalizability was achieved, but the unitarity is violated unless $a = 0$. This clearly shows that the renormalizability and unitarity exclude to each other. Although the a-term of providing massive graviton improves the ultraviolet divergence, it induces ghost excitations which jeopardizes the unitarity. In this sense, a first test for the quantum gravity is to require the unitarity, which means that there are no tachyon and ghost in its particle contents.

In this direction, recently, critical gravities have been an actively interesting subject because they were considered as toy models for quantum gravity [2, 3, 4, 5, 6, 7, 8]. At the critical point, a degeneracy take place and massive gravitons coincide with massless gravitons. Instead of massive gravitons, an equal amount of logarithmic modes appears in the critical gravity. According to the AdS/LCFT correspondence, one finds that a rank-2 logarithmic conformal field theory (LCFT) is dual to a critical gravity [9, 10, 11]. However, one has to resolve the non-unitarity issue of these log-gravity theories.

A polycritical gravity was, very recently, introduced to provide multiple critical points [12] which might be described by a higher-rank LCFT. The rank of the LCFT refers to the dimensionality of the Jordan cell. The LCFT dual to critical gravity has rank-2 and thus, an operator has one logarithmic partner. The LCFT dual to tricritical gravity has rank-3 and thus, an operator has two logarithmic partners. An odd-rank LCFT allows for a truncation to a unitary conformal field theory (CFT) [13]. A six-derivative gravity in three dimensions was treated as dual to a rank-3 LCFT [14], while a four-derivative critical gravity in four dimensions was considered as dual to a rank-3 LCFT [15]. Furthermore, it was shown that a consistent unitary truncation of polycritical gravity may be realized at the linearized level for odd rank [16]. On the other hand, a non-linear tricritical gravity of rank-3 in three and four dimensions was investigated in [17]. It is worth noting that a tricritical gravity was first mentioned in the six-derivative gravity in six dimensions [18].

Interestingly, one can construct a rank-3 parity odd theory in the context of four-derivative gravity known as three-dimensional generalized massive gravity (3DGMG) [19]. The 3DGMG is a combination of topologically massive gravity (TMG) [20] and new massive gravity (NMG) [21]. There exist two tricritical points in the 3DGMG parameter space whose dual theory is a rank-3 LCFT [22]. A truncation of the tricritical 3DGMG could be made by either imposing $Q_L = 0$ with Q_L the Abbott-Deser-Tekin charge or removing log-square boundary [14]. After truncation, a left-moving sector of CFT remains unitary.

At this stage, we would like to mention two related works on the 3DGMG. A log-square modes was found when choosing the Kerr-Schild metric [23] where its dual LCFT is not properly defined. Around the BTZ black hole, the authors [24] have confirmed the AdS/LCFT correspondence between the tricritical 3DGMG and a rank-3 finite temperature LCFT by computing quasinormal modes of graviton approximately. However, the 3DGMG is a toy model of tricritical gravities in three dimensions. Hence, it is very interesting to construct a four-dimensional generalized massive gravity (4DGMG).

In this work, we wish to construct the 4DGMG by combining quadratic gravity with the Chern-Simons modified term. Considering the AdS_4 vacuum solution together with the transverse-traceless gauge, we derive a lengthy linearized Einstein equation, which is not a compact form of the linearized equation as in the 3DGMG. Fortunately, taking the Kerr-Schild form of a perturbed metric tensor, the linearized tensor equation reduces to a single massive scalar equation on the AdS_4. At the tricritical points, we obtain a log-square solution which is similar to that found in the 3DGMG [23]. This is sharply compared to the three dimensional tricritical generalized massive gravity whose dual field theory is a rank-3 logarithmic conformal field theory.

2 General massive gravity in four dimensions

Before we proceed, we remind the reader that the 3DGMG is composed of NMG and TMG. Analogously, we introduce the quadratic gravity together with the Chern-Simons modified term [25, 26] in four dimensions as

$$S = \frac{1}{16\pi G}\int d^4x\sqrt{-g}\Big[R - 2\Lambda + \alpha R^2 + \beta R^{cd}R_{cd} + \frac{\theta}{4}{}^*RR\Big] \tag{2.1}$$

where Λ is a cosmological constant, α and β are parameters, while θ is a nondynamical field[1]. The inclusion of the nondynamical field comes from the fact that ${}^*RR$ is a total divergence [25, 26]. In this action, a term of ${}^*RR = {}^*R^a{}_b{}^{cd}R^b{}_{acd}$ represents the Pontryagin density, defined by

---

[1] We notice that in the dynamical Chern-Simons (DCS) modified gravity [26, 27, 28], the scalar field θ is treated as a dynamical field.

$$ {}^*R^a{}_b{}^{cd} = \frac{1}{2}\epsilon^{cdef}R^a{}_{bef}, \tag{2.2}$$

where ϵ^{cdef} denotes the four-dimensional Levi-Civita tensor. The last term corresponds to a parity-violating term of Chern-Simons modified gravity because of the parity operation $\hat{P}[{}^*RR] = -[{}^*RR]$. An important feature of this term shows that θ is realized as $\nabla_c\theta$ in the equation of motion, which may spoil the diffeomorphism [see Eq.(2.6)].

Varying for g_{ab} on the action (2.1) leads to the Einstein equation

$$R_{ab} - \frac{1}{2}g_{ab}R + \Lambda g_{ab} + E_{ab} + C_{ab} = 0, \tag{2.3}$$

where E_{ab} and the four-dimensional Cotton tensor C_{ab} are given by

$$\begin{aligned} E_{ab} &= \alpha\left(2RR_{ab} - \frac{1}{2}g_{ab}R^2 + 2g_{ab}\nabla^2R - \nabla_a\nabla_bR\right) \\ &\quad +\beta\left(g_{ab}\nabla^2R - \nabla_a\nabla_bR + \nabla^2R_{ab} - \frac{1}{2}g_{ab}\nabla^2R + 2R_{acbd}R^{cd} - \frac{1}{2}g_{ab}R_{cd}R^{cd}\right), \end{aligned} \tag{2.4}$$

$$C_{ab} = \nabla_c\,\theta\,\epsilon^{cde}{}_{(a}\nabla_{|e|}R_{b)d} + \frac{1}{2}\nabla_c\nabla_d\,\theta\,\epsilon_{(b}{}^{cef}R^d{}_{a)ef}. \tag{2.5}$$

Here, the Cotton tensor C_{ab} is a traceless and symmetric tensor. It is worth noting that applying ∇_a to (2.3), we get a remaining term as the Bianchi identity

$$\nabla^aC_{ab} = -\left[\frac{\nabla_b\theta}{8}\right]{}^*RR, \tag{2.6}$$

which should be zero when choosing a proper background solution. If not, the diffeomorphism-symmetry breaking is being realized from the fact that the covariant divergence of the Cotton tensor is non-zero, which contrasts to the case in three dimensions. Hence, a consistency condition implies that ${}^*RR = 0$ for $\nabla_b\theta \neq 0$ because the other case of $\nabla_b\theta = 0 (C_{ab} = 0)$ reduces to the quadratic gravity. In this sense, the diffeomorphism-symmetry is restored dynamically for the case of ${}^*RR = 0$ when choosing the AdS$_4$-vacuum in the equation of motion, even though it may occur at the action level. We note that the same conclusion is found when θ is considered as a Lagrange multiplier.

On the other hand, it turns out that Eq.(2.3) has an AdS$_4$ solution in which the Riemann tensor, Ricci tensor and Ricci scalar are given by

$$\bar{R}_{abcd} = \frac{\Lambda}{3}(\bar{g}_{ac}\bar{g}_{bd} - \bar{g}_{ad}\bar{g}_{bc}), \quad \bar{R}_{ab} = \Lambda\bar{g}_{ab}, \quad \bar{R} = 4\Lambda. \tag{2.7}$$

Here "overbar" denotes the background AdS$_4$-metric $\bar{g}_{ab}$. In order to obtain the linearized equation, we introduce the perturbation around the the background metric given by

$$g_{ab} = \bar{g}_{ab} + h_{ab}. \tag{2.8}$$

In this case, the linearized equation to (2.3) can be written as

$$\delta R_{ab}(h) - \frac{1}{2} g_{ab} \delta R(h) - \Lambda h_{ab} + \delta E_{ab} + \delta C_{ab}(h) = 0, \tag{2.9}$$

where the linearized tensor $\delta R_{ab}(h)$, $\delta R(h)$, $\delta E_{ab}(h)$ [3] and $\delta C_{ab}(h)$ [29] are given by

$$\begin{aligned}
\delta R_{ab}(h) &= \frac{1}{2}\left(\bar{\nabla}^c \bar{\nabla}_a h_{bc} + \bar{\nabla}^c \bar{\nabla}_b h_{ac} - \bar{\nabla}^2 h_{ab} - \bar{\nabla}_a \bar{\nabla}_b h\right), \\
\delta R(h) &= \bar{\nabla}^a \bar{\nabla}^b h_{ab} - \bar{\nabla}^2 h - \Lambda h, \\
\delta E_{ab}(h) &= 4\Lambda\left(2\alpha + \frac{1}{3}\beta\right)\delta G_{ab} + (2\alpha + \beta)\left(g_{ab}\bar{\nabla}^2 - \bar{\nabla}_a \bar{\nabla}_b + \Lambda g_{ab}\right)\delta R \\
&\quad + \beta\left(\bar{\nabla}^2 \delta G_{ab} - \frac{2\Lambda}{3} g_{ab} \delta R\right), \\
\delta C_{ab}(h) &= \Bigg[\frac{1}{2} v_c\ \epsilon^{cde}{}_a\left(\bar{\nabla}_e \delta R_{bd} - \Lambda \bar{\nabla}_e h_{bd}\right) + \frac{1}{8} v_{cd}\ \epsilon_b{}^{cef}\Big(\bar{\nabla}_e \bar{\nabla}_f h^d{}_a + \bar{\nabla}_e \bar{\nabla}_a h^d{}_f \\
&\quad - \bar{\nabla}_e \bar{\nabla}^d h_{af} - \bar{\nabla}_f \bar{\nabla}_e h^d{}_a - \bar{\nabla}_f \bar{\nabla}_a h^d{}_e + \bar{\nabla}_f \bar{\nabla}^d h_{ae}\Big)\Bigg] + \Big[a \leftrightarrow b\Big]
\end{aligned} \tag{2.10}$$

with

$$\delta G_{ab} = \delta R_{ab} - \frac{1}{2}\delta R \bar{g}_{ab} - \Lambda h_{ab}, \quad v_c = \bar{\nabla}_c \theta, \quad v_{cd} = \bar{\nabla}_c \bar{\nabla}_d \theta. \tag{2.11}$$

Imposing the transverse gauge as

$$\bar{\nabla}^a h_{ab} = \bar{\nabla}_b h \tag{2.12}$$

and substituting (2.12) into Eq.(2.9), the trace part of the perturbation equation (2.9) becomes

$$\Lambda\Big(h - 2(3\alpha + \beta)\bar{\nabla}^2 h\Big) = 0, \tag{2.13}$$

which results in $\bar{g}^{ab}\delta C_{ab} = 0$. In order to eliminate a massive spin-0 mode, we set [3]

$$3\alpha + \beta = 0. \tag{2.14}$$

In this case, Eq.(2.13) yields $h = 0$ which implies that we may have the transverse traceless (TT) gauge

$$\bar{\nabla}^a h_{ab} = 0, \quad h = 0. \tag{2.15}$$

Substitution of (2.14) and (2.15) into Eq.(2.9) leads to

$$\frac{3\alpha}{2}\left(\bar{\nabla}^2 - \frac{2\Lambda}{3}\right)\left(\bar{\nabla}^2 - \frac{4\Lambda}{3} - \frac{1}{3\alpha}\right) h_{ab} + \delta C_{ab}(h) = 0. \tag{2.16}$$

Here the linearized Cotton tensor $\delta C_{ab}(h)$ is given by

$$\begin{aligned}\delta C_{ab}(h) \;\; = \;\; & \Big[-\frac{1}{4} v_c\ \epsilon^{cde}{}_a \bar{\nabla}_e \bar{\nabla}^2 h_{bd} + \frac{\Lambda}{6} v_c\ \epsilon^{cde}{}_a \bar{\nabla}_e h_{bd} + \frac{1}{4} v_{cd}\ \epsilon_b{}^{cef}\Big(\bar{\nabla}_e \bar{\nabla}_a h^d{}_f - \bar{\nabla}_e \bar{\nabla}^d h_{af}\Big)\Big] \\ & + \Big[a \leftrightarrow b\Big]. \end{aligned} \tag{2.17}$$

At this stage, we point out that it is difficult to combine δC_{ab} with the first term in Eq.(2.16) without any choice of θ. It is, however, known that for a particular choice of θ [29, 30], one can manipulate δC_{ab} so that the perturbation equation (2.16) can be factored out. This could be achieved by introducing the Poincare coordinates (u, v, x, y) for the AdS$_4$:

$$\theta = k\frac{x}{y}, \quad \bar{g}_{ab} \equiv \phi^{-2}\eta_{ab} = \frac{\ell^2}{y^2}\eta_{ab}, \tag{2.18}$$

where k has the dimension of $[\text{mass}]^{-2}$, ℓ is the AdS$_4$ curvature radius ($\Lambda = -3/\ell^2$) and η_{ab} is the flat metric tensor defined as

$$\eta_{ab}dx^a dx^b = 2dudv + dx^2 + dy^2. \tag{2.19}$$

Plugging Eq.(2.18) into Eq.(2.16), one arrives at

$$\left(\bar{\nabla}^2 - \frac{2}{3}\Lambda\right)\left[- 3\alpha\Big(\bar{\nabla}^2 - \frac{4}{3}\Lambda - \frac{1}{3\alpha}\Big) h_{ab} + v_c\ \epsilon^{cde}{}_{(a}\bar{\nabla}_{|e|} h_{b)d}\right] = 0. \tag{2.20}$$

Alternatively, it leads to

$$\left[- 3\alpha\Big(\bar{\nabla}^2 - \frac{4}{3}\Lambda - \frac{1}{3\alpha}\Big)\delta^{a'}_{(a}\delta^d_{b)} + \delta^{a'}_{(a} v_{|c|}\ \epsilon^{cde}{}_{b)}\bar{\nabla}_e\right]\left(\bar{\nabla}^2 - \frac{2}{3}\Lambda\right) h_{a'd} = 0 \tag{2.21}$$

because two operations in Eq.(2.20) commute to each other. Here, v_c is given by

$$v_c = k\left(0, 0, \frac{1}{y}, -\frac{x}{y^2}\right) \tag{2.22}$$

which generates the mass of graviton. In this case, v_c is not a constant vector but a vector field.

3 AdS wave as perturbation

It is a formidable task to solve (2.21) directly because the four dimensional Cotton term induces a complicated expression upon choosing v_c as in (2.22). In order to solve Eq.(2.21), we take the Kerr-Schild form as an AdS$_4$-wave solution

$$h_{ab} = 2\varphi \lambda_a \lambda_b, \tag{3.1}$$

where $\lambda_a = (1,0,0,0)$ is a null ($\lambda^2 = 0$), and geodesic vector and a scalar field $\varphi = \varphi(u,v,x,y)$. The TT gauge condition (2.15) implies that one may restrict φ to $\varphi(u,x,y)$ by requiring the condition of $\lambda_a \bar{\nabla}^a \varphi = 0$. Substituting $h_{ab} = 2\varphi\lambda_a\lambda_b$ into the tensor Eq.(2.20) leads to the scalar equation for φ

$$\begin{aligned} \lambda_{a'}\lambda_d \Big[-3\alpha \Big(\bar{\nabla}^2 + \frac{4}{\phi}\partial^f \phi \, \partial_f - \frac{1}{3\alpha}\Big) \delta^{a'}_{(a}\delta^{d}_{b)} + \delta^{a'}_{(a} v_{|c|} \, \epsilon^{cde}{}_{b)} \Big(\frac{\partial_e \phi}{\phi} + \bar{\nabla}_e \Big) \Big] & \\ \times \Big[\bar{\nabla}^2 + \frac{2}{3}\Lambda + \frac{4}{\phi}\partial^f \phi \, \partial_f \Big] \varphi = 0. & \end{aligned} \tag{3.2}$$

Introducing the separation of variables $\varphi(u,x,y) = U(u)X(x)Y(y)$ and taking into account λ_a, ϕ (2.18) and v_c (2.22), Eq.(3.2) reduces to

$$\begin{aligned} \Bigg[3\alpha \Big(y^2(\partial_y^2 + \partial_x^2) + 2y\partial_y - \frac{\ell^2}{3\alpha}\Big) + k\Big(x\partial_x + y\partial_y + 1\Big) \Bigg] & \\ \times \Bigg(y^2(\partial_y^2 + \partial_x^2) + 2y\partial_y - 2 \Bigg) XY = 0. & \end{aligned} \tag{3.3}$$

We remark that for the AdS$_4$-wave, the second parenthesis in Eq.(3.3) is consistent with the linearized equation of the massless scalar, which corresponds to the second parenthesis of massless tensor in Eq.(2.21).

On the other hand, as was suggested in the 3DGMG [19], the first bracket in Eq.(3.3) is related to the massive equation which consists of two mass parameters m_1 and m_2. In order to obtain two massive equations from the first bracket in Eq.(3.3), we introduce the four first-order operators,

$$D^{m_i} = y\partial_y + m_i, \quad D^{-1} = y\partial_y - 1, \quad D^{+2} = y\partial_y + 2, \tag{3.4}$$

where $i = 1, 2$. Using these mutually commuting operators, Eq.(3.3) can be expressed compactly as

$$D^{m_1}D^{m_2}D^{-1}D^{+2}\, Y(y) = 0 \to (y\partial_y + m_1)(y\partial_y + m_2)(y\partial_y - 1)(y\partial_y + 2)\, Y(y) = 0. \tag{3.5}$$

In deriving this, we assume that $X(x)$ =constant. Comparing Eq.(3.5) with Eq.(3.3), we find m_1 and m_2 ($\alpha \neq 0$)

$$m_1 = \frac{1}{2}\left(1+\frac{k}{3\alpha}+\sqrt{\frac{k^2}{9\alpha^2}-\frac{2k}{3\alpha}+1+\frac{4\ell^2}{3\alpha}}\right), \tag{3.6}$$

$$m_2 = \frac{1}{2}\left(1+\frac{k}{3\alpha}-\sqrt{\frac{k^2}{9\alpha^2}-\frac{2k}{3\alpha}+1+\frac{4\ell^2}{3\alpha}}\right). \tag{3.7}$$

For $\alpha = 0$, being the Chern-Simons modified gravity [29], we have a third-order perturbation equation from Eq.(3.3) instead of the fourth-order equation (3.5). In this case, a single mass parameter m can be determined as

$$m_1 = m = 1-\frac{\ell^2}{k},\quad m_2 = 0 \quad \text{or} \quad m_2 = 1-\frac{\ell^2}{k},\quad m_1 = 0. \tag{3.8}$$

4 Tricritical and critical points

Now we wish to obtain tricritical points which can be read off from Eq. (3.5). For this purpose, we focus on two points as follows:

$$\text{point 1}: \quad k = 9\alpha = -3\ell^2 \;\Rightarrow\; m_1 = m_2 = 2 \tag{4.1}$$

$$\text{point 2}: \quad k = -9\alpha = \frac{3}{4}\ell^2 \;\Rightarrow\; m_1 = m_2 = -1. \tag{4.2}$$

For each point, equation (3.5) reduces to

$$D^{-1}D^{+2}D^{+2}D^{+2}\,Y(y) = 0 \;\Rightarrow\; (y\partial_y+2)^3(y\partial_y-1)\,Y(y) = 0 \tag{4.3}$$

$$D^{-1}D^{-1}D^{-1}D^{+2}\,Y(y) = 0 \;\Rightarrow\; (y\partial_y-1)^3(y\partial_y+2)\,Y(y) = 0, \tag{4.4}$$

respectively. It is important to note that at the point 1, the operators D^{m_1} and D^{m_2} degenerate with D^{+2} [Eq.(4.3)], while at the point 2, they degenerate with D^{-1} [Eq.(4.4)]. We remind the reader that these correspond to two tricritical points, which implies that there exists three-fold degeneracy.

On the other hand, it is well known that in the quadratic gravity [31], obtained from taking the $k = 0$ limit in Eq.(3.3), the critical point is given by $\alpha = \ell^2/6 = -1/(2\Lambda)$. In this case, one checks that Eq.(3.3) reduces to

$$D^{-1}D^{+2}D^{-1}D^{+2}\,Y(y) = 0 \;\Rightarrow\; (y\partial_y-1)^2(y\partial_y+2)^2\,Y(y) = 0. \tag{4.5}$$



Figure 1: The tricritical and critical points on the k vs α graph for $\ell^2 = 1$. The tricritical points 1 and 2 correspond to $(\alpha,\ k) = (-1/3,\ -3)$ and $(-1/12,\ 3/4)$, while three critical points $3-5$ to $(1/6,\ 0)$, $(0,\ 1/2)$, $(0,\ -1)$, respectively. In addition, the point $(0,\ 0)$ denotes the Einstein gravity.

In the Chern-Simons modified gravity [29], the $\alpha = 0$ limit in Eq. (3.3), we obtain two critical points of $k = \ell^2/2$ and $k = -\ell^2$ from Eq.(3.8). For each k, Eq. (3.3) becomes the third-order equation as

$$D^{-1}D^{-1}D^{+2}\,Y(y) = 0 \;\;\Rightarrow\;\; (y\partial_y + 2)(y\partial_y - 1)^2\,Y(y) = 0, \quad (k = \ell^2/2) \tag{4.6}$$

$$D^{-1}D^{+2}D^{+2}\,Y(y) = 0 \;\;\Rightarrow\;\; (y\partial_y - 1)(y\partial_y + 2)^2\,Y(y) = 0. \quad (k = -\ell^2) \tag{4.7}$$

For $k = \ell^2/2$, two-fold degeneracy emerges at $m = m_1 = -1$, whereas there exists two-fold degeneracy at $m = m_2 = 2$ for $k = -\ell^2$.

In what follows, we find the solution to Eqs.(4.3)-(4.7) for each point:

(i) $m_1 = m_2 = 2$ (point 1 : tricritical)

$$\begin{aligned}\varphi(u,y) &= U(u)Y(y) \\ &= c_1(u)y + \frac{1}{y^2}\Big[c_2(u) + c_3(u)\ln(y) + c_4(u)\ln^2(y)\Big],\end{aligned} \tag{4.8}$$

(ii) $m_1 = m_2 = -1$ (point 2 : tricritical)

$$\begin{aligned}
\varphi(u,y) &= U(u)Y(y) \\
&= c_5(u)\frac{1}{y^2}+y\Big[c_6(u)+c_7(u)\ln(y)+c_8(u)\ln^2(y)\Big],
\end{aligned} \tag{4.9}$$

(iii) $m_1=2$, $m_2=-1$ (point 3 : critical)

$$\begin{aligned}
\varphi(u,y) &= U(u)Y(y) \\
&= d_1(u)y+d_2(u)\frac{1}{y^2}+d_3(u)y\ln(y)+d_4(u)\frac{\ln(y)}{y^2},
\end{aligned} \tag{4.10}$$

(iv) $m_1=-1$, $m_2=0$ (point 4 : critical)

$$\begin{aligned}
\varphi(u,y) &= U(u)Y(y) \\
&= d_5(u)y+d_6(u)\frac{1}{y^2}+d_7(u)y\ln(y),
\end{aligned} \tag{4.11}$$

(v) $m_1=0$, $m_2=2$ (point 5 : critical)

$$\begin{aligned}
\varphi(u,y) &= U(u)Y(y) \\
&= d_8(u)y+d_9(u)\frac{1}{y^2}+d_{10}(u)\frac{\ln(y)}{y^2},
\end{aligned} \tag{4.12}$$

where $c_{1\sim 8}$ and $d_{1\sim 10}$ are undetermined functions of u.

From (i)-(v), we observe that at tricritical points 1 and 2, there exists the log-square term, as was shown in the tricritical 3DGMG [23], while at the critical points of $3-5$, the solutions have the log terms. In particular, the combination of the solution (4.11) and (4.12) yields the solution (4.10) in the quadratic gravity [31] which provides the fourth-order perturbation equation at the critical point.

In summary, Fig.1 shows two tricritical points and three critical points in the 4DGMG, which are depicted in the $(\alpha,\ k)$ plane.

5 Discussions

We have constructed a generalized massive gravity by combining quadratic gravity with the Chern-Simons modified term in four dimensions. The four-dimensional generalized massive gravity is similar to the three-dimensional massive gravity. Considering the AdS$_4$ vacuum solution, we have derived the linearized Einstein equation, which is not a compact form in compared to that of the three-dimensional generalized massive gravity. For the Kerr-Schild form of a perturbed metric tensor, however, the linearized tensor equation reduces to a massive scalar equation. At the tricritical points, we obtain a log-square wave solution whose dual field theory is not properly defined. This is compared clearly to the three-dimensional tricritical generalized massive gravity whose dual field theory is a rank-3 logarithmic conformal field theory.

A crucial difference between the 4DGMG and 3DGMG is that the former has the Chern-Simons modified term of $\frac{\theta}{4}\,{}^*RR$ with θ a function of the spacetimes, while the latter has the topologically massive term $\frac{1}{2\mu}\epsilon\cdot[\Gamma\partial\Gamma+\frac{2}{3}\Gamma\Gamma\Gamma]$, even though two terms are parity-violating term. The presence of θ in the 4DGMG prevents us from having its dual LCFT at the critical point because it is a nondynamical field but not a constant like $1/\mu$ in the 3DGMG. Most of all, choosing $\theta = kx/y$ makes the linearized equation complicated on the AdS$_4$ spacetimes, which implies that the equation can not be written by a compact form, as was shown in the AdS$_3$ background:

$$(\mathcal{D}^L\mathcal{D}^R\mathcal{D}^{M+}\mathcal{D}^{M-}h)_{\mu\nu}=0, \tag{5.1}$$

where $h_{\mu\nu}$ is the perturbation of the metric around the AdS$_3$ spacetimes and the differential operators $\mathcal{D}^{L,\ R}$, $\mathcal{D}^{M+,\ M-}$ are given in [14, 19, 24].

Therefore, the parity-odd tricritical gravity theory in the 4DGMG is very difficult to have its dual higher rank LCFT except the tricritical 3DGMG.

Acknowledgments

This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MEST) through the Center for Quantum Spacetime (CQUeST) of Sogang University with grant number 2005-0049409. Y. Myung was partly supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MEST) (No.2011-0027293).

References

[1] K. S. Stelle, Phys. Rev. D **16**, 953 (1977).

[2] W. Li, W. Song and A. Strominger, JHEP **0804**, 082 (2008) [arXiv:0801.4566 [hep-th]].

[3] H. Lu and C. N. Pope, Phys. Rev. Lett. **106**, 181302 (2011) [arXiv:1101.1971 [hep-th]].

[4] S. Deser, H. Liu, H. Lu, C. N. Pope, T. C. Sisman and B. Tekin, Phys. Rev. D **83**, 061502 (2011) [arXiv:1101.4009 [hep-th]].

[5] M. Porrati and M. M. Roberts, Phys. Rev. D **84**, 024013 (2011) [arXiv:1104.0674 [hep-th]].

[6] M. Alishahiha and R. Fareghbal, Phys. Rev. D **83**, 084052 (2011) [arXiv:1101.5891 [hep-th]].

[7] E. A. Bergshoeff, O. Hohm, J. Rosseel and P. K. Townsend, Phys. Rev. D **83**, 104038 (2011) [arXiv:1102.4091 [hep-th]].

[8] H. Lu, C. N. Pope, E. Sezgin and L. Wulff, JHEP **1110**, 131 (2011) [arXiv:1107.2480 [hep-th]].

[9] D. Grumiller and N. Johansson, JHEP **0807**, 134 (2008) [arXiv:0805.2610 [hep-th]].

[10] Y. S. Myung, Phys. Lett. B **670**, 220 (2008) [arXiv:0808.1942 [hep-th]].

[11] A. Maloney, W. Song and A. Strominger, Phys. Rev. D **81**, 064007 (2010) [arXiv:0903.4573 [hep-th]].

[12] T. Nutma, Phys. Rev. D **85**, 124040 (2012) [arXiv:1203.5338 [hep-th]].

[13] E. A. Bergshoeff, S. de Haan, W. Merbis, M. Porrati and J. Rosseel, JHEP **1204**, 134 (2012) [arXiv:1201.0449 [hep-th]].

[14] E. A. Bergshoeff, S. de Haan, W. Merbis, J. Rosseel and T. Zojer, arXiv:1206.3089 [hep-th].

[15] N. Johansson, A. Naseh and T. Zojer, arXiv:1205.5804 [hep-th].

[16] A. Kleinschmidt, T. Nutma and A. Virmani, arXiv:1206.7095 [hep-th].

[17] L. Apolo and M. Porrati, arXiv:1206.5231 [hep-th].

[18] H. Lu, Y. Pang and C. N. Pope, Phys. Rev. D **84**, 064001 (2011) [arXiv:1106.4657 [hep-th]].

[19] Y. Liu and Y. -W. Sun, Phys. Rev. D **79**, 126001 (2009) [arXiv:0904.0403 [hep-th]].

[20] S. Deser, R. Jackiw and S. Templeton, Ann. Phys. (N.Y.) **140**, 372 (1982); **185**, 406(E) (1988); **281**, 409 (2000).

[21] E. A. Bergshoeff, O. Hohm and P. K. Townsend, Phys. Rev. Lett. **102**, 201301 (2009) [arXiv:0901.1766 [hep-th]].

[22] D. Grumiller, N. Johansson and T. Zojer, JHEP **1101**, 090 (2011) [arXiv:1010.4449 [hep-th]].

[23] E. Ayon-Beato, G. Giribet and M. Hassaine, JHEP **0905**, 029 (2009) [arXiv:0904.0668 [hep-th]].

[24] Y. -W. Kim, Y. S. Myung and Y. -J. Park, arXiv:1207.3149 [hep-th].

[25] R. Jackiw and S. Y. Pi, Phys. Rev. D **68**, 104012 (2003) [gr-qc/0308071].

[26] S. Alexander and N. Yunes, Phys. Rept. **480**, 1 (2009) [arXiv:0907.2562 [hep-th]].

[27] V. Cardoso and L. Gualtieri, Phys. Rev. D **80**, 064008 (2009) [Erratum-ibid. D **81**, 089903 (2010)] [arXiv:0907.5008 [gr-qc]].

[28] T. Moon and Y. S. Myung, Phys. Rev. D **84**, 104029 (2011) [arXiv:1109.2719 [gr-qc]], and references therein.

[29] T. Moon and Y. S. Myung, Eur. Phys. J. C **71**, 1796 (2011) [arXiv:1108.2612 [hep-th]].

[30] E. A.-Beato, G. Giribet, and M. Hassaine, Phys. Rev. **D83**, 104033 (2011) [arXiv:1103.0742 [hep-th]].

[31] I. Gullu, M. Gurses, T. C. Sisman and B. Tekin, Phys. Rev. D **83**, 084015 (2011) [arXiv:1102.1921 [hep-th]].